

***Acanthopanax senticosus* polysaccharide suppressing proliferation and metastasis of the human non-small cell lung cancer NCI-H520 cells is associated with Wnt/ β -catenin signaling**

D. SUN^{1,*}, J. CHEN², H. HU³, S. LIN³, L. JIN¹, L. LUO⁴, X. YAN³, C. ZHANG⁵

¹Institute of Life Sciences, Wenzhou University, Wenzhou, China; ²Sichuan Provincial Medical Center of Mental Health & Sichuan Provincial People's Hospital, Sichuan, China; ³School of Pharmaceutical Sciences, Wenzhou Medical University, Wenzhou, China; ⁴Bioengineering College, Chongqing University, Chongqing, China; ⁵The Third Affiliated Hospital of Wenzhou Medical University, Wenzhou, China

*Correspondence: sunda_dsd@163.com

Received September 13, 2018 / Accepted January 15, 2019

Lung cancer (LC) has the highest mortality of all tumors. Non-small cell lung cancer (NSCLC) accounts for about 80% of all LC. *Acanthopanax senticosus* polysaccharide (ASPS) is extracted from the root of *Acanthopanax senticosus* (AS). Herein, we examined the effect and molecular mechanism of ASPS on NSCLC. The proliferation, invasion and migration of NCI-H520 cells were detected by cell counting kit-8 (CCK-8), transwell assay and wound healing assay, respectively. The epithelial-mesenchymal transition (EMT) and Wnt/ β -catenin pathway-related factors were evaluated using quantitative real-time PCR (qRT-PCR) and western blot assay. Our results showed that ASPS significantly decreased the proliferation of cells at 24 and 48 h. Moreover, ASPS markedly repressed the invasion and migration capacities of cells in a concentration-dependent manner. Besides, ASPS obviously downregulated the levels of matrix metalloproteinase-2 (MMP-2), MMP-9, fibronectin 1 (FN1), vimentin, wnt3a, phosphorylated-glycogen synthase kinase 3 β (p-GSK3 β) and cyclin D1, whereas E-cadherin level was upregulated. The level of GSK3 β was not changed within the different groups. ASPS conspicuously inhibited the abilities of proliferation and metastasis in human non-small cell lung cancer cell line NCI-H520 possibly by suppressing Wnt/ β -catenin pathway mediated-EMT.

Key words: *Acanthopanax senticosus* polysaccharide, proliferation, metastasis, Wnt/ β -catenin pathway

Lung cancer (LC) has the highest mortality of all tumors and about 1.5 million people die of LC each year worldwide. According to the histologic characteristic, LC is divided into two types: small cell lung cancer (SCLC) and non-small cell lung cancer (NSCLC) [1]. The proportion of NSCLC patients in newly diagnosed LC patients is up to 85% per year and the 5-year survival rate is less than 15% [2]. Although chest screening, low dose CT and other early screening methods are widely used in the early detection of NSCLC, the most patients are in middle and late stage when diagnosed, thereby losing the chance of surgical cure [3, 4]. At the present, treatment of NSCLC mainly includes chemotherapy, molecular targeted therapy and immunotherapy [4]. However, these treatments have some side effects and the patient's survival rate is still very low [5, 6]. Hence, we need to examine the specific pathogenesis of NSCLC and find a safe and effective treatment.

It is well known that the widespread use of Chinese herbal medicine in the treatment of various tumors is a hot spot of research [7–9]. Researchers have proved that the antitumor activity of Chinese herbal medicine is mainly derived from the plant extracts, including polysaccharides, saponins, steroids, etc. [10–12]. In recent years, the extensive biological activity of plant polysaccharides has been gradually recognized [13]. The antitumor research of polysaccharide has aroused wide attention of scholars both at home and abroad [14, 15].

Acanthopanax senticosus (AS) belongs to Chinese herbal medicine [16]. *Acanthopanax senticosus* polysaccharide (ASPS) is extracted from the root of AS and is one of the main active components of AS [17]. The bioactivities of ASPS include immunomodulatory, antitumor, antioxidant, anti-radiation and anti-inflammatory [18–21]. Meanwhile, it has the advantages of low toxicity, so it has a great potential in

clinical application [22, 23]. Nevertheless, the role of ASPS in NSCLC is not yet known.

The development of tumor is the evolutionary process of multifactor, multigene and multistage gradual accumulation. Cell signal transduction pathway plays a pivotal role in its development, invasion and metastasis [24, 25]. As we know, the Wnt/ β -catenin signal transduction pathway plays an important regulatory role in cell differentiation, proliferation, survival and apoptosis, as well as cell carcinogenesis, tumor invasion and other pathological processes [26]. Wnt/ β -catenin pathway is closely related to the tumors and the abnormal activation of the pathway is involved in the pathogenesis of a variety of human cancers, such as nasopharyngeal carcinoma, gastric cancer, liver cancer, breast cancer, prostate cancer, etc. [27–32].

At the present study, we explored the effect of ASPS on the proliferation and metastasis of human non-small cell lung cancer line NCI-H520 and further studied whether the potential mechanism was the Wnt/ β -catenin pathway involvement.

Materials and methods

Drug treatment. *Acanthopanax senticosus* polysaccharide (ASPS) was extracted from the cortex of *Acanthopanax senticosus*. The powder of *Acanthopanax senticosus* polysaccharide was purchased from Shaanxi Ciyuan Biotechnology Co., Ltd. The color was brown and the purity was more than 80%. The powder was dissolved in serum-free Roswell Park Memorial Institute-1640 (RPMI-1640) (Biosun, Shanghai, China) and filtered through 0.22 μ m membrane (Shenghua; Haining, Jiaxing, China). The solution of ASPS was diluted to 10, 20, 40, 80, 160 and 320 mg/ml in RPMI-1640.

Cell culture. Human non-small cell lung cancer cell line NCI-H520 was obtained from Shanghai enzyme linked Biotechnology Co., Ltd. Cells were incubated in RPMI-1640 supplemented with 10% Fetal Bovine Serum (FBS; Lonsera, Shanghai, China) and penicillin-streptomycin solution

(Leagene, Beijing, China) in 37°C incubator (S-G80A-II; Sheyanyiqi, Shanghai, China) with 95% relative humidity and 5% CO₂.

Cell Counting Kit-8 (CCK-8) assay. CCK-8 (Yeasen, Shanghai, China) was used to assess the cell proliferation as manufacturer's instructions recommend. Firstly, cells were inoculated into 96-well plates at the density of 1.5×10³ cells/well and cultured in incubator for 24 h. Next day, cells were treated with PBS and different concentration of ASPS (0, 10, 20, 40, 80, 160 and 320 mg/ml) for 12, 24 and 48 h, respectively. Then, CCK-8 solution was dripped into each well and the plate was transferred into the incubator for 4 h. Finally, OD value at 450 nm was detected by HBS-1096B microplate reader (Detie, Nanjing, China).

Transwell assay. Transwell assay was carried out to evaluate cell invasion according to the manufacturer's instructions. Briefly, cells were inoculated into 6-well plates at the density of 2×10⁴ cells/well and cultured in incubator for 24 h. Next day, cells were treated with PBS (control) and different concentration of ASPS: 40 (ASPS1), 80 (ASPS2) and 160 (ASPS3) mg/ml for 48 h, respectively. BD matrigel (Solarbio, Shanghai, China) was filled in the upper chamber of transwell at 37°C for 30 min. Then, transwell was placed into the culture plate. RPMI-1640 was added into the upper chamber of transwell for 25 min and sucked away. 10% FBS was dropped into the lower chamber of the transwell. Subsequently, cell suspension was cultured in the upper chamber of transwell for 24 h. Cells were stained with 0.1% crystal violet (Zhongxin chemtech; Tianjin, China) for 20 min and washed with PBS for 3 times. Finally, cells were observed and photographed using inverted microscope (DYS-339; Dianying, Shanghai, China).

Wound healing assay. Wound healing test was performed to analyze cell migration following a standardized method. Firstly, cells were inoculated into 6-well plates at the density of 2×10⁴ cells/well and cultured in incubator for 24 h. Next day, cells were scratched about 500 μ m width by 1 mL pipette tip and washed with RPMI-1640 for 2 times. Then, cells were treated with PBS (control) and different concentration of ASPS at 40 (ASPS1), 80 (ASPS2) and 160 (ASPS3) mg/ml for 0, 24 and 48 h, respectively. Finally, cells were observed and photographed using inverted microscope.

Quantitative real-time PCR (qRT-PCR) assay. Total RNA of was collected by TRIeasy™ Total RNA Extraction Reagent (Yeasen, Shanghai, China). cDNA was synthesized by ABScript II cDNA first strand synthesis kit (ABclonal, Beijing, China). The reaction reagents were as follows: 1 μ g RNA, 2 μ l d(T)₂₃VN, 10 μ l ABScript II reaction mix, 2 μ l ABScript II enzyme mix, 1 μ l dNTP_s, nuclease-free H₂O to a total volume of 20 μ l. Reaction conditions were 80°C for 5 min. Then, cDNA was amplified by SYBR Premix Taq™ II kit (Takara, Beijing, China). The reaction reagents were as follows: 25 μ l SYBR Green Mix, 4 μ l cDNA, 1 μ l forward/reverse primer, nuclease-free H₂O to a total volume of 50 μ l. Reaction conditions were as followed: predegeneration: 95°C

Table 1. Sequences of the primers.

Primer name	Sequence (5'-3')	Product size (bp)
MMP-2-Forward	CAGCCCTGCAAGTTTCCATT	210
MMP-2-Reverse	GTTGCCCCAGGAAAGTGAAGG	
MMP-9-Forward	GAGACTCTACCCCAGGACG	238
MMP-9-Reverse	GAAAGTGAAGGGGAAGACGC	
FN1-Forward	TGGCACTGATGAAGAACCCT	224
FN1-Reverse	GGGAAACTGTGTAGGGGTCA	
Vimentin-Forward	AATAAGATCCTGCTGGCCGA	225
Vimentin-Reverse	GGTGTTCGCTTCCTCTC	
E-cadherin-Forward	ACGCATTGCCACATACACTC	217
E-cadherin-Reverse	GGTGTTCACATCATCGTC	
β -actin-Forward	GGGAAATCGTGCGTGACATT	219
β -actin-Reverse	AGGTAGTTTCGTGGATGCCA	

for 5 min, (denaturation: 95 °C for 15 s; annealing: 62 °C for 35 s) for 40 cycles, extending: 75 °C for 30 s. The primer sequence was listed in the Table 1. β -actin was regarded as internal control. The formula $2^{-\Delta\Delta CT}$ was used to compare the quantification.

Western blot assay. Total protein was lysed with high RIPA lysis buffer (Leagene, Beijing, China). Protein was separated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and bound to polyvinylidene difluoride (PVDF) membrane (Thermo Fisher Scientific, Shanghai, China). Then, membrane was sealed by 5% skim milk at room temperature for 1.5 h and incubated with anti-matrix metalloproteinase-2 (MMP-2) (R&D, IC903G-100UG, 1:1000), anti-MMP-9 (Abcam, EP1254, 1:500), anti-fibronectin 1 (FN1) (Abcam, ab32419, 1:800), anti-vimentin (R&D, AF2105, 1:700), anti-E-cadherin (R&D, MAB1838, 1:1000), anti-wnt3a (Abcam, ab28472, 1:800), anti- β -catenin (R&D, AF1329, 1:1000), anti-phosphorylated-glycogen synthase kinase 3 β (p-GSK3 β) (Abcam, ab75814, 1:1000), anti-GSK3 β (Abcam, ab32391, 1:1000), anti-cyclin D1 (R&D,

MAB4314, 1:900) and anti- β -actin (Abcam, ab13772, 1:800) at 4 °C refrigerator overnight. Next day, membrane was incubated with the secondary antibodies (Rabbit anti-mouse IgG, CST, #58802, 1:7000; Goat anti-rabbit IgG, Beyotime, A0423, 1:8000; Donkey anti-goat IgG, Beyotime, A0181, 1:8000) at 37 °C for 60 min. The protein was exposed using the ECL chromogenic solution (Biodragon, Beijing, China).

Statistical analysis. All experiments were carried out at least three times. Experimental data was presented as mean \pm standard deviation (SD) using Microsoft Excel. One-way ANOVA was performed to evaluate the differences between groups. A $p < 0.05$ was treated as a significant difference.

Results

ASPS decreased the proliferation of NCI-H520 cells.

The effect of ASPS on the proliferation of NCI-H520 cells was measured by CCK-8. The data showed that there was no obvious change in cell proliferation, when cells were administrated to ASPS at 10, 20, 40, 80, 160 and 320 mg/ml for




Figure 1. The effect of ASPS on cell viabilities of NCI-H520 cell and human pulmonary alveolar epithelial cell (PAEpiC). (A) NCI-H520 cells were administrated to PBS (0) and different concentrations of ASPS: 0, 10, 20, 40, 80, 160 and 320 mg/ml for 12, 24 and 48 h, respectively. (B) PAEpiC cells were administrated to PBS (0) and different concentrations of ASPS at 0, 40, 80, 160 and 320 mg/ml for 24 and 48 h, respectively. CCK-8 was used to evaluate the cell proliferation. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. control (PBS).

12 h. Besides, ASPs markedly inhibited cell proliferation for 24 and 48 h and 80 mg/ml ASPs significantly inhibited cell proliferation compared to control (0 mg/ml) at 24 h. The IC₅₀ of ASPs was 160 mg/ml at 48 h. When the cells were treated with 320 mg/ml for 48 h, the rate of cell proliferation was less than 50% (Figure 1A). Moreover, study also demonstrated that 160 mg/ml ASPs treatment for 24 and 48 h had no significant inhibitory effect on human pulmonary alveolar epithelial cell viability (Figure 1B). Therefore, 40, 80 and 160 mg/ml ASPs treatment for 48 h were worth further investigating and were determined as the later experimental condition.

ASPs repressed the invasion of NCI-H520 cells. The effect of ASPs on the invasion of NCI-H520 cells was detected by transwell assay. Our results showed that the number of cell invasion was drastically reduced, when cells were administrated to ASPs. As quantitative analysis of cell invasiveness showed, the rates of cell invasion were 65%, 45% and 30% in ASPs1, ASPs2 and ASPs3 groups, respectively. In comparison with control group, the invasiveness of cells was decreased by 35%, 55% and 70% in ASPs1, ASPs2 and ASPs3 groups, respectively (Figures 2A–B).

ASPs suppressed the migration of NCI-H520 cells. The effect of ASPs on the migration of NCI-H520 cells was analyzed by wound healing test. Our data showed that the number of cell migration was dramatically reduced, when

the cells were administrated to ASPs. As quantitative analysis of cell migration showed, cell migration rates were 80%, 56% and 33% in ASPs1, ASPs2 and ASPs3 groups for 24 h, respectively. Cell migration rates were 60%, 50% and 25% in ASPs1, ASPs2 and ASPs3 groups for 48 h, respectively. The rates of cell migration were lessened by 20%, 44% and 67% in ASPs1, ASPs2 and ASPs3 groups for 24 h, respectively, compared with the control group. The rates of cell migration were reduced by 40%, 50% and 75% for 48 h, respectively, compared to the control group (Figures 3A–B).

ASPs suppressed the metastasis of NCI-H520 cells by regulating EMT-related factors. qRT-PCR and western blot assays were used to assess EMT-related factors. The qRT-PCR data revealed that the RNA levels of MMP-2, MMP-9, FN1 and vimentin were obviously declined and E-cadherin RNA level was enhanced in ASPs groups in dose-dependent manner, relative to the control group (Figure 4A). In addition, western blot data showed that the expression trend of protein was the same as that of RNA (Figure 4B).

ASPs suppressed the proliferation and metastasis of NCI-H520 cells by downregulating Wnt/ β -catenin pathway. The molecular mechanism of ASPs on the proliferation and metastasis of NCI-H520 cells was examined by western blot assay. As western blot results showed that as compared with control group, ASPs significantly downreg-




Figure 2. ASPs repressed the invasion of NCI-H520 cells. A) Cells were treated with PBS (control) and different concentrations of ASPs: 40 (ASPs1), 80 (ASPs2) and 160 (ASPs3) mg/ml for 48 h, respectively. Transwell assay was performed to detect cell invasion. B) Quantitative analysis of cell invasiveness was carried out with GraphPad prism 7. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. control.




Figure 3. ASPS suppressed the migration of NCI-H520 cells. A) Cells were treated with PBS (control) and different concentrations of ASPS: 40 (ASPS1), 80 (ASPS2) and 160 (ASPS3) mg/ml for 0, 24 and 48 h, respectively. Wound healing assay was used to explore cell migration. B) Quantitative analysis of cell migration rate was carried out with GraphPad prism 7. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. control.

ulated the protein levels of wnt3a, β -catenin, cyclin D1 in a concentration-dependent manner. Meanwhile, the phosphorylation level of GSK3 β was attenuated in ASPS groups in dose-dependent manner. Moreover, there was no significant change in the expression of GSK3 β protein between different groups (Figure 5).

Discussion

For decades, accumulating evidence testifies that plant polysaccharides have obvious anticancer activity [14, 15, 33].

Han et al. have found that *Sargassum henslowianum* polysaccharide can treat gastric cancer by improving the immunomodulatory function [14]. Ayeka PA *et al.* have confirmed that licorice polysaccharide inhibits the growth of tumor, thereby improves the health of mice [33]. Therefore, we inferred that ASPS had the effect of anti-NSCLC. As expected, ASPS distinctly repressed the proliferation of NCI-H520 cells in concentration-dependent manner at 24 and 48 h.

Cancer metastasis is the most dangerous stage in the process of tumorigenesis and evolution and is the most important cause of cancer death, as well a key factor




Figure 4. ASPs suppressed the metastasis of NCI-H520 cells by regulating EMT-related factors. A) RNA levels of MMP-2, MMP-9, FN1, vimentin and E-cadherin were analyzed by qRT-PCR assay. B) The protein levels of MMP-2, MMP-9, FN1, vimentin and E-cadherin were measured by western blot assay and normalized to β -actin expression. Gray value was detected and counted by quality one. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. control.




Figure 5. ASPs suppressed the proliferation and metastasis of NCI-H520 cells by downregulating Wnt/ β -catenin pathway. The protein levels of Wnt3a, β -catenin, Cyclin D1, p-GSK3 β and GSK3 β were evaluated by western blot assay. β -actin was regarded as internal control. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. control.

affecting the effect of tumor treatment [34]. Recent research has demonstrated that *Hizikia fusiformis* polysaccharide represses the invasion ability of human fibrosarcoma cells [35]. Furthermore, it has been found that *Acorus calamus* L. polysaccharide reduces the metastasis of lung cancer cells [36]. Similar to previous studies, our data showed that ASPS markedly reduced the invasion and migration capacities of NCI-H520 cells in a dose-dependent manner.

Metastasis of tumor cells involves the regulation of epithelial-mesenchymal transition (EMT) process-related molecules. EMT-related factors include MMPs, FN, vimentin, E-cadherin, and so on [37–39]. A study has proved that FBXO11 facilitates the capacities of proliferation and metastasis in gastric cancer cells through inducing EMT process [37]. Another research confirms that SPARC reduces the growth, invasion and migration of T-cell non-Hodgkin's lymphoma cells by inhibiting EMT process [40]. In addition, accumulating evidence substantiates that when the metastasis of the tumor cells is suppressed, the EMT process is blocked, accompanied by a decrease in the expression levels of MMP-2, MMP-9, vimentin and FN and an increase in E-cadherin expression [38, 39]. In this study, we observed that ASPS repressed the proliferation and metastasis of NCI-H520 cells by blocking EMT process, which downregulated levels of MMP-2, MMP-9, vimentin and FN1 and upregulated E-cadherin levels.

A large number of studies have shown that genes and drugs inhibit the growth and metastasis of various cancer cells by regulating the Wnt/ β -catenin pathway [27–29, 31]. Moreover, Wnt/ β -catenin pathway can induce EMT transformation by inhibiting GSK3 β -mediated phosphorylation and the degradation of β -catenin and further represses downstream target genes, such as cyclin D1 [31, 41]. A previous report indicated that transmembrane 4 L6 family proteins silencing depressed the metastasis of NSCLC via downregulating Wnt/ β -catenin and EMT signaling [42]. As expected, our results illustrated that ASPS significantly lessened metastasis of NCI-H520 cells via Wnt/ β -catenin signaling pathway mediated-EMT. Though, validation experiments were also required, it was the limitation of the current study and we are planning to use antagonists or agonists to validate the possible signaling pathway.

All in all, our research testified that ASPS significantly repressed the proliferation, invasion and migration abilities of human non-small cell lung cancer NCI-H520 cells through suppressing Wnt/ β -catenin pathway mediated-EMT. Based on the results of this study, ASPS may be developed as a drug for the treatment of NSCLC in the future.

Acknowledgements: This work was supported by Wenzhou Science & Technology Innovation Project in Water Pollution Control & Manage [Grant number: W20170010]; Zhejiang Pharmaceutical Top Priority Academic Open Fund [Grant number: 201720]. The analyzed data sets generated during the study are available from the corresponding author on reasonable request.

References

- [1] JEMAL A, BRAY F, CENTER MM, FERLAY J, WARD E et al. Global cancer statistics. *CA Cancer J Clin* 2011; 61: 69–90. <https://doi.org/10.3322/caac.20107>
- [2] HUANG YT, HEIST RS, CHIRIEAC LR, LIN X, SKAUG V et al. Genome-wide analysis of survival in early-stage non-small-cell lung cancer. *J Clin Oncol* 2009; 27: 2660–2667. <https://doi.org/10.1200/JCO.2008.18.7906>
- [3] CHEN Z, WANG J, BAI Y, WANG S, YIN X et al. The associations of TERT-CLPTM1L variants and TERT mRNA expression with the prognosis of early stage non-small cell lung cancer. *Cancer Gene Ther* 2017; 24: 20–27. <https://doi.org/10.1038/cgt.2016.74>
- [4] RUSSO AE, PRIOLO D, ANTONELLI G, LIBRA M, MCCUBREY JA et al. Bevacizumab in the treatment of NSCLC: patient selection and perspectives. *Lung Cancer (Auckl)* 2017; 8: 259–269. <https://doi.org/10.2147/LCTT.S110306>
- [5] BEGUM S, HAYASHI M, OGAWA T, JABBOURE FJ, BRAIT M et al. An integrated genome-wide approach to discover deregulated microRNAs in non-small cell lung cancer: Clinical significance of miR-23b-3p deregulation. *Sci Rep* 2015; 5: 13236. <https://doi.org/10.1038/srep13236>
- [6] RECK M, POPAT S, REINMUTH N, DE RUYSSCHER D, KERR KM et al. Metastatic non-small-cell lung cancer (NSCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2014; 25 Suppl 3: iii27–39. <https://doi.org/10.1093/annonc/mdu199>
- [7] PAN J, XU Y, SONG H, ZHOU X, YAO Z et al. Extracts of Zuo Jin Wan, a traditional Chinese medicine, phenocopies 5-HTR1D antagonist in attenuating Wnt/beta-catenin signaling in colorectal cancer cells. *BMC Complement Altern Med* 2017; 17: 506. <https://doi.org/10.1186/s12906-017-2006-7>
- [8] WANG Q, WANG Q, WANG SF, JIAO LJ, ZHANG RX et al. Oral Chinese herbal medicine as maintenance treatment after chemotherapy for advanced non-small-cell lung cancer: a systematic review and meta-analysis. *Curr Oncol* 2017; 24: e269–e276. <https://doi.org/10.3747/co.24.3561>
- [9] WANG S, WU X, TAN M, GONG J, TAN W et al. Fighting fire with fire: poisonous Chinese herbal medicine for cancer therapy. *J Ethnopharmacol* 2012; 140: 33–45. <https://doi.org/10.1016/j.jep.2011.12.041>
- [10] PRADO S, FERREIRA GF, HARAZONO Y, SHIGA TM, RAZ A et al. Ripening-induced chemical modifications of papaya pectin inhibit cancer cell proliferation. *Sci Rep* 2017; 7: 16564. <https://doi.org/10.1038/s41598-017-16709-3>
- [11] QIAN Y, HAN QH, LIU D, TU PE, ZENG KW et al. [Antitumor target identification and molecular mechanism study of total saponins from *Albizia julibrissin*]. *Zhongguo Zhong Yao Za Zhi* 2017; 42: 3661–3665. <https://doi.org/10.19540/j.cnki.cjcmm.2017.0144>
- [12] WANG JN, ZHANG Y, SONG LY, PENG YB, LI WJ. [Chemical constituents from *Pteris dispar* and their antitumor activity in vitro]. *Zhongguo Zhong Yao Za Zhi* 2017; 42: 4159–4164. <https://doi.org/10.19540/j.cnki.cjcmm.20170905.002>

- [13] TAN X, ZHOU X, CHEN HG. [Structure-activity relationship of plant polysaccharides]. *Zhongguo Zhong Yao Za Zhi* 2017; 42: 4104–4109. <https://doi.org/10.19540/j.cnki.cjcm.20170928.016>
- [14] HAN M, SUN P, LI Y, WU G, NIE J. Structural characterization of a polysaccharide from *Sargassum henslowianum*, and its immunomodulatory effect on gastric cancer rat. *Int J Biol Macromol* 2018; 108: 1120–1127. <https://doi.org/10.1016/j.ijbiomac.2017.12.109>
- [15] KUMAR SSD, MAHESH A, ANTONIRAJ MG, RATHORE HS, HOURELD NN et al. Cellular imaging and folate receptor targeting delivery of gum kondagogu capped gold nanoparticles in cancer cells. *Int J Biol Macromol* 2018; 109: 220–230. <https://doi.org/10.1016/j.ijbiomac.2017.12.069>
- [16] HAN Y, ZHANG A, SUN H, ZHANG Y, MENG X et al. High-throughput ultra high performance liquid chromatography combined with mass spectrometry approach for the rapid analysis and characterization of multiple constituents of the fruit of *Acanthopanax senticosus* (Rupr. et Maxim.) Harms. *J Sep Sci* 2017; 40: 2178–2187. <https://doi.org/10.1002/jssc.201601445>
- [17] HAN J, LI JH, BAI G, SHEN GS1, CHEN J et al. *Acanthopanax senticosus* polysaccharides-induced intestinal tight junction injury alleviation via inhibition of NF-kappaB/MLCK pathway in a mouse endotoxemia model. *World J Gastroenterol* 2017; 23: 2175–2184. <https://doi.org/10.3748/wjg.v23.i12.2175>
- [18] FEI XJ, ZHU LL, XIA LM, PENG WB, WANG Q. *Acanthopanax senticosus* attenuates inflammation in lipopolysaccharide-induced acute lung injury by inhibiting the NF-kappaB pathway. *Genet Mol Res* 2014; 13: 10537–10544. <https://doi.org/10.4238/2014.December.12.16>
- [19] HAN J, LIU L, YU N, CHEN J, LIU B et al. Polysaccharides from *Acanthopanax senticosus* enhances intestinal integrity through inhibiting TLR4/NF-kappaB signaling pathways in lipopolysaccharide-challenged mice. *Anim Sci J* 2016; 87: 1011–1018. <https://doi.org/10.1111/asj.12528>
- [20] WANG H, SUN B, ZHANG Z, CHEN J, HAO Q et al: Effects of *Acanthopanax senticosus* polysaccharide on the proliferation, apoptosis and cell cycle in human HepG2 cells. *Pharmazie* 2016; 71: 201–204. <https://doi.org/10.1691/ph.2016.5807>
- [21] XIE SS. [Immunoregulatory effect of polysaccharide of *Acanthopanax senticosus* (PAS). I. Immunological mechanism of PAS against cancer]. *Zhonghua Zhong Liu Za Zhi* 1989; 11: 338–340.
- [22] BANG JS, CHUNG YH, CHUNG SJ, LEE HS, SONG EH et al. Clinical effect of a polysaccharide-rich extract of *Acanthopanax senticosus* on alcohol hangover. *Pharmazie* 2015; 70: 269–273. <https://doi.org/10.1691/ph.2015.4786>
- [23] HUANG L, ZHAO H, HUANG B, ZHENG C, PENG W et al. *Acanthopanax senticosus*: review of botany, chemistry and pharmacology. *Pharmazie* 2011; 66: 83–97. <https://doi.org/10.1691/ph.2011.0744>
- [24] HAHNFELDT P, PANIGRAHY D, FOLKMAN J, HLATKY L. Tumor development under angiogenic signaling: a dynamical theory of tumor growth, treatment response, and postvascular dormancy. *Cancer Res* 1999; 59: 4770–4775.
- [25] TURASHVILI G, BROGI E. Tumor Heterogeneity in Breast Cancer. *Front Med (Lausanne)* 2017; 4: 227. <https://doi.org/10.3389/fmed.2017.00227>
- [26] LOGAN CY, NUSSE R. The Wnt signaling pathway in development and disease. *Annu Rev Cell Dev Biol* 2004; 20: 781–810. <https://doi.org/10.1146/annurev-cell-bio.20.010403.113126>
- [27] FU Q, GAO Y, YANG F, MAO T, SUN Z et al. Suppression of microRNA-454 impedes the proliferation and invasion of prostate cancer cells by promoting N-myc downstream-regulated gene 2 and inhibiting WNT/beta-catenin signaling. *Biomed Pharmacother* 2018; 97: 120–127. <https://doi.org/10.1016/j.biopha.2017.10.115>
- [28] HUANG J, HE Y, MCLEOD HL, XIE Y, XIAO D et al. miR-302b inhibits tumorigenesis by targeting EphA2 via Wnt/beta-catenin/EMT signaling cascade in gastric cancer. *BMC Cancer* 2017; 17: 886. <https://doi.org/10.1186/s12885-017-3875-3>
- [29] KRISHNAMURTHY N, KURZROCK R. Targeting the Wnt/beta-catenin pathway in cancer: Update on effectors and inhibitors. *Cancer Treat Rev* 2018; 62: 50–60. <https://doi.org/10.1016/j.ctrv.2017.11.002>
- [30] LI X, LI M, CHEN J, DAI H, WANG L et al. SAMMSON drives the self-renewal of liver tumor initiating cells through EZH2-dependent Wnt/beta-catenin activation. *Oncotarget* 2017; 8: 103785–103796. <https://doi.org/10.18632/oncotarget.21792>
- [31] WANG W, WEN Q, LUO J, CHU S, CHEN L et al. Suppression Of beta-catenin Nuclear Translocation By CGP57380 Decelerates Poor Progression And Potentiates Radiation-Induced Apoptosis in Nasopharyngeal Carcinoma. *Theranostics* 2017; 7: 2134–2149. <https://doi.org/10.7150/thno.17665>
- [32] YANG J, HAN F, LIU W, CHEN H, HAO X et al. ALX4, an epigenetically down regulated tumor suppressor, inhibits breast cancer progression by interfering Wnt/beta-catenin pathway. *J Exp Clin Cancer Res* 2017; 36: 170. <https://doi.org/10.1186/s13046-017-0643-9>
- [33] AYEKA PA, BIAN Y, GITHAIGA PM, ZHAO Y. The immunomodulatory activities of licorice polysaccharides (*Glycyrrhiza uralensis* Fisch.) in CT 26 tumor-bearing mice. *BMC Complement Altern Med* 2017; 17: 536. <https://doi.org/10.1186/s12906-017-2030-7>
- [34] KU KH, PARK SJ, KIM JH, KWON HJ, CHANG HK et al. [Gastric Cancer Recurrence in 12 Years after Surgical Resection]. *Korean J Gastroenterol* 2017; 70: 296–300. <https://doi.org/10.4166/kjg.2017.70.6.296>
- [35] LEE SG, KARADENIZ F, OH JH, YU GH, KONG CS. Inhibitory Effect of *Hizikia fusiformis* Solvent-Partitioned Fractions on Invasion and MMP Activity of HT1080 Human Fibrosarcoma Cells. *Prev Nutr Food Sci* 2017; 22: 184–190. <https://doi.org/10.3746/pnf.2017.22.3.184>
- [36] LOPATINA KA, SAFONOVA EA, NEVSKAYA KV, STAKHEEVA MN, GUREV AM et al. Effect of *Acorus calamus* L. Polysaccharide on CD274 and CD326 Expression by Lewis Lung Carcinoma Cells in Mice. *Bull Exp Biol Med* 2017; 164: 102–105. <https://doi.org/10.1007/s10517-017-3934-4>

- [37] SUN C, TAO Y, GAO Y, XIA Y, LIU Y et al. F-box protein 11 promotes the growth and metastasis of gastric cancer via PI3K/AKT pathway-mediated EMT. *Biomed Pharmacother* 2018; 98: 416–423. <https://doi.org/10.1016/j.biopha.2017.12.088>
- [38] WU A, LI M, MAI Z, LI S, YANG Z. [CK2alpha Regulates the Metastases and Migration of Lung Adenocarcinoma A549 Cell Line through PI3K/Akt/GSK-3beta Signal Pathway]. *Zhongguo Fei Ai Za Zhi* 2017; 20: 233–238. <https://doi.org/10.3779/j.issn.1009-3419.2017.04.11>
- [39] XU S, WANG T, GUAN ZR, ZHANG C, CHEN Y et al. FBXO2, a novel marker for metastasis in human gastric cancer. *Biochem Biophys Res Commun* 2018; 495: 2158–2164. <https://doi.org/10.1016/j.bbrc.2017.12.097>
- [40] YAN J, ZHANG J, ZHANG X, LI X, LI L et al. SPARC is down-regulated by DNA methylation and functions as a tumor suppressor in T-cell lymphoma. *Exp Cell Res* 2018; 364: 125–132. <https://doi.org/10.1016/j.yexcr.2017.12.022>
- [41] WU Y, TRAN T, DWABE S, SARKISSYAN M, KIM J et al. A83-01 inhibits TGF-beta-induced upregulation of Wnt3 and epithelial to mesenchymal transition in HER2-overexpressing breast cancer cells. *Breast Cancer Res Treat* 2017; 163: 449–460. <https://doi.org/10.1007/s10549-017-4211-y>
- [42] CHOI SI, KIM SY, LEE JH, KIM JY, CHO EW et al. Osteopontin production by TM4SF4 signaling drives a positive feedback autocrine loop with the STAT3 pathway to maintain cancer stem cell-like properties in lung cancer cells. *Oncotarget* 2017; 8: 101284–101297. <https://doi.org/10.18632/oncotarget.21021>